

Block Designs for Partially Diallel Cross

Kuey Chung Choi

Division of Mathematics, Computer Science and Statistics, Chosun University,

375 Seosuk-Dong Dong-Gu,

Kwangju, Korea.

kjchoi@mail.chosun.ac.kr

Sudhir Gupta

Division of Statistics, Northern Illinois University,

DeKalb, IL 60115, U.S.A.

sudhir@math.niu.edu

Young Nam Son

The Research Institute of Statistics, Chosun University,

375 Seosuk-Dong Dong-Gu,

Kwangju, Korea.

syn2000@netian.com

1. Introduction

Consider a block design D_b for a diallel cross experiment involving n_c distinct crosses laid out in b blocks of k crosses each, with each cross replicated r_c times. Following Singh and Hinkelmann (1995), the model for the data is assumed to be $Y = \mu 1_n + \Delta_1 g + \Delta_2 \beta + \varepsilon$ where the symbols have their usual meaning. The purpose of this paper is to define partially balanced (partial) diallel cross designs and to give some methods of constructing them.

Definition. A diallel cross design D_b will be called an m -associate class partially balanced diallel cross (PBDC) design with parameters $\{p, n_c, b, r_c, k, \alpha_1, \alpha_2, \dots, \alpha_m\}$ if the following holds for a given pair of lines β and γ that are i th associates: $k\lambda_{c(\beta,\gamma)} - \lambda_{b(\beta,\gamma)} = \alpha_i$ where $\lambda_{b(\beta,\gamma)}$ and $\lambda_{c(\beta,\gamma)}$ are the numbers of concurrences of the lines β and γ in designs D_b and D_c respectively, and α_i is a constant independent of the pair of i th associates chosen, $i = 1, 2, \dots, m$.

2. Two general methods of construction

We now present two widely applicable methods of constructing PBDC designs. Let D_1 be an m -associate class PBIB design with parameters $v = p, b = b_1, r = r_1, k = 2, \lambda_1, \lambda_2, \dots, \lambda_m$ such that $\lambda_i = 0$ for $i(\neq s) = 1, 2, \dots, m$, and $\lambda_s = 1$, where $s \in \{1, 2, \dots, m\}$.

Theorem 1. The existence of an m -associate class PBIB design D_1 with parameters $p, b_1, r_1, k_1 = 2, \lambda_s = 1, \lambda_i = 0, i(\neq s) = 1, 2, \dots, m$, and the existence of a BIB design D_2 with parameters $v = b_1, b_2, r_2, k_2, \lambda$ implies the existence of an m -associate class PBDC design with parameters $\{p, n_c = b_1, b = b_2, r_c = r_2, k = k_2, \alpha_s = \lambda(b_1 - r_1^2), \alpha_i = -r_1^2\lambda, i(\neq s) = 1, 2, \dots, m\}$.

The following gives a method of construction using α -resolvable PBIB designs.

Theorem 2. The existence of an α -resolvable m -associate class PBIB design D_1 with parameters $p, b_1, r_1, k = 2, \lambda_i = 0$ or $1, i = 1, 2, \dots, m$ implies the existence of an m -associate class PBDC design with parameters $\{p, n_c = b_1, b = r_1/\alpha, k = \alpha p/2, \alpha_i = k\lambda_i - \alpha r_1, i = 1, 2, \dots, m\}$.

REFERENCES

- Choi, K.C. and Gupta, S. (2000). On constructions of optimal complete diallel crosses. *Utilitas Math.*, **58**, 153-160.
- Das, A. and Ghosh, D.K. (1999). Balanced incomplete block diallel cross designs. *Statistics and Applications*, **1**, 1-16.
- Das, A., Dean, A.M. and Gupta, S. (1998). On optimality of some partial diallel cross designs, *Sankhya*, **B 60**, 511-524.
- Das, A., Dey, A. and Dean, A.M. (1998). Optimal designs for diallel cross experiments, *Statist. & Prob. Letters*, **36**, 427-436.
- Dey, A. and Midha, C.K. (1996). Optimal block designs for diallel crosses, *Biometrika*, **83**, 484-489.
- Gupta, S. and Kageyama, S. (1994). Optimal complete diallel crosses, *Biometrika*, **81**, 420-424.
- S. Gupta and M. Singh (1989). Analysis of PBIB designs using association matrices. *Metrika* **36**, 1-6.
- Prasad, R., Gupta, V.K. and R. Srivastava. (1999). Universally optimal block designs for diallel crosses, *Statistics and Applications*, **1**, 35-52.
- Singh, M. and Hinkelmann, K. (1995). Partial diallel crosses in incomplete blocks, *Biometrics*, **51**, 1302-1314.

RESUME

Kuey Chung Choi

Professor and Chairman, Research Institute of Statistics

Division of Mathematics, Computer Science and Statistics, Chosun University

375 Seosuk-Dong, Kwangju, 501-759, Republic of Korea

e-mail: kjchoi@mail.chosun.ac.kr

Phone : +82-62-230-6622